## **Table Notes**

## **Coronary Artery Bypass Graft (CABG)**

**Total Number of Cases** represents all inpatient hospitalizations, after exclusions, for patients 18 years and older who underwent a CABG (without a valve procedure).

**Mortality** represents patients who died during the hospital stay.

**Readmission** represents patients who were readmitted to a Pennsylvania acute care hospital within 7, 30 and 90 days of the discharge date of the original hospitalization. Out-of-state residents were excluded because readmission data was not available for patients readmitted to a non-Pennsylvania hospital. Planned readmissions were not counted.

## Extended Postoperative Length of Stay

represents patients whose length of stay in the hospital following a CABG procedure was significantly longer than expected, after accounting for patient risk.

Average Hospital Charge represents the entire length of stay and is trimmed and case-mix adjusted. Professional fees were not included. In almost all cases, hospitals typically receive actual payments from private insurers or government payers that are considerably less than the listed charge. Coronary artery bypass graft (CABG) surgery is used to treat a blockage in a coronary artery by creating an alternate pathway for the blood to flow in order to reach vital heart muscle. CABG is typically recommended for severe or complex blockages that are not treatable by other methods.

## **Understanding the Symbols**

The symbols displayed in this report represent a comparison of actual *mortality*, *readmission* and *extended postoperative length of stay* rates to what is expected, after accounting for patient risk.

Using readmission as an example:

- O Rate was significantly lower than expected. Fewer patients were readmitted than could be attributed to patient risk and random variation.
- Rate was not significantly different than expected. The number of patients who were readmitted was within the range anticipated based on patient risk and random variation.
- Rate was significantly higher than expected. More patients were readmitted than could be attributed to patient risk and random variation.

See About the Report section or Technical Notes for further details.